

# Drug Facts

---

## Active ingredients (in each 59-mL delivered dose)

### Purpose

Monobasic sodium phosphate 9.5 g....	Saline laxative
Dibasic sodium phosphate 3.5 g....	Saline laxative

---

**Use** for relief of occasional constipation

---

## Warnings

For rectal use only

---

### Dosage Warning:

Using more than one enema in 24 hours can be harmful. Do not use without asking a doctor:

- for more than 3 days
  - if your child has kidney disease
  - if your child has heart problems
  - if your child is dehydrated
- 

**Ask a doctor before using this product if the child**

- is on a sodium-restricted diet
- 

**Ask a doctor before using any laxative if your child has**

- abdominal pain, nausea, or vomiting
  - a sudden change in bowel habits lasting more than 2 weeks
  - already used a laxative for more than 1 week
- 

**Stop use and ask a doctor if the child has**

- rectal bleeding
- no bowel movement or no liquid comes out of the child's rectum after 30 minutes
- symptoms of dehydration [feeling thirsty, dizziness, vomiting, urinating less often than normal]

These symptoms may indicate a serious condition

---

**Keep out of reach of children to prevent accidental ingestion.** If swallowed, get medical help or contact a Poison Control Center right away. ►

# Drug Facts (continued)

## Directions

Single Daily Dosage (per 24 hours)

**Do not use more per day unless directed by a doctor.**

**See Warnings.**

**Do not use if child is taking another sodium phosphates product.**

children 5 to 11 years	1 bottle per day or as directed by doctor
------------------------	---

children 2 to under 5 years	one-half bottle per day (see below)
-----------------------------	-------------------------------------

children under 2 years	DO NOT USE
------------------------	------------

## One-half bottle preparation:

Unscrew cap and remove 2 tablespoons of liquid with a measuring spoon. Replace cap and follow "How to use this enema" directions.

## Other information

- each 59-mL delivered dose contains: sodium 2.2g
- additional liquids by mouth are recommended while taking this product
- this product usually produces a bowel movement in 1-5 minutes
- **BOTTLE SEALED FOR SAFETY. IF BOTTLE WRAPPER WITH FLEET® EMBLEM IS TORN, DAMAGED OR MISSING, DO NOT USE**
- for complete professional use warnings and precautions, consult the Reference Safety Information on our website

## Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

**Questions? 1-866-255-6960**

**www.pedia-lax.com**