

Drug Facts

Active ingredient Purpose (in each tablet)

Magnesium hydroxide 400 mg.....Saline laxative

Uses

- for relief of occasional constipation
- this product generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before using any laxative if your child has

- kidney disease
- a magnesium-restricted diet
- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if your child

- has rectal bleeding
- does not have a bowel movement within 6 hours of taking this product

These symptoms may be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

Use dosage chart for proper dosing. Doses may be taken as a single daily dose or in divided doses. Drink a full glass (8 ounces) of liquid with each dose.

Age	Starting Dose	Maximum dose per day
children 6 to under 12 years	3-6 tablets	6 tablets
children 2 to under 6 years	1-3 tablets	3 tablets
children under 2 years	ask a doctor	

Other information

- each tablet contains: magnesium 170 mg
- CHILD RESISTANT CAP
- Tablet may not appear as shown
- The top of the bottle is sealed with foil for your safety. Do not use if foil imprinted "SEALED for YOUR PROTECTION" is broken or missing.

Inactive ingredients

colloidal silicon dioxide, FD&C red #40 aluminum lake, flavor, magnesium stearate, maltodextrin, mannitol, sorbitol, stearic acid, sucralose

Questions? 1-866-255-6960 or www.pedia-lax.com