

Drug Facts

Active ingredient **Purpose**
(in each 2.3-mL average delivered dose)

Glycerin 2.3 g.....Hyperosmotic laxative

Uses

- for relief of occasional constipation
- this product generally produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

May cause rectal discomfort or a burning sensation.

Ask a doctor before using any laxative if the child has

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if the child has

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions Single daily dosage

children 2 to under 6 years	1 suppository or as directed by a doctor
children under 2 years	consult a doctor

Drug Facts (continued)

Directions (continued)

Left-side position: Place child on left side with knees bent and arms resting comfortably.

Knee-chest position: Have child kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.

Administering the laxative: **CAUTION: REMOVE ORANGE PROTECTIVE SHIELD BEFORE INSERTING.**

Hold unit upright, grasping bulb of unit with fingers. Grasp orange protective shield with other hand, pull gently to remove. With steady pressure, gently insert tip into rectum with a slight side-to-side movement, with tip pointing toward navel. **Discontinue use if resistance is encountered. Forcing the tip can result in injury.** Insertion may be easier if child receiving suppository bears down as if having a bowel movement. This helps relax the muscles around the anus. Squeeze the bulb until nearly all liquid is expelled. While continuing to squeeze the bulb, remove tip from rectum and discard unit. It is not necessary to empty unit completely. The unit contains more than the amount of liquid needed for effective use. A small amount of liquid will remain in the unit after squeezing.

Other information

- carton sealed for safety. If seal with Fleet emblem on bottom flap or top flap is broken or missing, do not use

Inactive ingredient purified water

Questions? 1-866-255-6960 or www.pedia-lax.com