

Drug Facts

Active ingredient (in each tablespoon)

Docusate sodium 50 mg Stool softener

Purpose

Uses

- for the prevention of dry, hard stools
- for relief of occasional constipation
- this product generally produces bowel movement in 12-72 hours

Warnings

Drug interaction precaution: Do not give this product to your child if your child is presently taking mineral oil, unless directed by a doctor

Ask a doctor before using any laxative if your child has

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if your child has

- rectal bleeding
- no bowel movement within 72 hours of using this product

These symptoms may be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Use dosage chart for proper dosing. Doses may be taken as a single daily dose or in divided doses. Doses must be given in a 6-8 oz glass of milk or juice, to prevent throat irritation.

Age	Starting Dose	Maximum dose per day
children 2 to under 12 years	1-3 tablespoons	3 tablespoons
children under 2 years	ask a doctor	

Drug Facts (continued)

Other information

- each tablespoon contains: sodium 13 mg
- CHILD RESISTANT CAP
- The top of the bottle is sealed with foil for your safety. Do not use if foil imprinted "SEALED for YOUR PROTECTION" is broken or missing.

Inactive ingredients

citric acid, edetate disodium, FD&C red #3, flavor, methylparaben, polyethylene glycol, povidone, propylene glycol, propylparaben, sodium citrate, sorbitol, sucralose, water, xanthan gum, xylitol

Questions? 1-866-255-6960
or www.pedia-lax.com

© C.B. Fleet Company, Inc. 2007
The package design is a trademark of
C.B. Fleet Company, Inc.
Lynchburg, VA 24502 U.S.A.

Fleet
Laboratories, Inc.