

Drug Facts

Active ingredient (in each suppository)

Purpose

Glycerin 1g.....Hyperosmotic laxative

Uses

- for relief of occasional constipation
- this product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

May cause rectal discomfort or a burning sensation.

Ask a doctor before using any laxative if child has

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop use and consult a doctor if child has

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

Keep out of reach of children to prevent accidental ingestion.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Single daily dosage (per 24 hours)

children 2 to under 6 years	1 suppository or as directed by a doctor
children under 2 years	ask a doctor

Insert suppository well up into rectum.

To produce laxative action suppository is designed to dissolve only partially, which may not be noticeable.

Other information

- Store container tightly closed.
- Keep away from excessive heat.
- Mouth of jar sealed for safety.
If foil imprinted with Green Fleet® emblem is missing or torn, do not use.

Inactive ingredients

edetate disodium, purified water, sodium hydroxide, stearic acid

Questions? 1-866-255-6960 or
www.pedia-lax.com